

Variable, but not free-weight, resistance exercise potentiates jump performance following a task-specific warm-up.

í de la companya de l	
Journal:	Scandinavian Journal of Medicine and Science in Sports
Manuscript ID	SJMSS-0-716-18.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	07-Nov-2018
Complete List of Authors:	Mina, Minas; University of Derby, School of Sport, Outdoor and Exercise Science Blazevich, Anthony; Edith Cowan University, Centre for Exercise and Sports Science Research Tsatalas, Themistoklis; University of Thessaly, Department of Physical Education and Sport Science Giakas, Giannis; University of Thessaly, Department of Physical Education and Sport Science Seitz, Laurent; Edith Cowan University, Centre for Exercise and Sports Science Research Kay, Anthony; The University of Northampton, Sport, Exercise & Life Sciences
Keywords:	elastic bands, PAP, conditioning activity, explosive strength, kinetics, kinematics

SCHOLARONE[™] Manuscripts

1		
2 3 4	1	TITLE
5 6	2	Variable, but not free-weight, resistance back squat exercise potentiates jump performance
7 8 0	3	following a comprehensive task-specific warm-up
9 10 11	4	AUTHORS
12 13	5	Minas A. Mina ¹ , Anthony J. Blazevich ² , Themistoklis Tsatalas ³ , Giannis Giakas ³ , Laurent B.
14 15 16	6	Seitz ² & Anthony D. Kay ⁴
17 18	7	
19 20	8	INSTITUTIONAL AFFILIATIONS
21 22 23	9	¹ Sport, Outdoor and Exercise Science, University of Derby, Derby, United Kingdom
23 24 25	10	² Centre for Exercise and Sports Science Research, School of Medical & Health Sciences, Edith
26 27	11	Cowan University, Joondalup, Western Australia
28 29 20	12	³ Department of Physical Education & Sport Science, University of Thessaly, Greece
30 31 32	13	⁴ Sport, Exercise & Life Sciences, University of Northampton, Northampton, United Kingdom
33 34	14	
35 36	15	RUNNING HEAD
37 38 39	16	Elastic band use improves jump performance
40 41	17	
42 43	18	CORRESPONDING AUTHOR
44 45 46	19	Minas A. Mina ¹
40 47 48	20	Sport, Outdoor and Exercise Science
49 50	21	University of Derby
51 52 53	22	Kedleston Road, Derby
55 54 55	23	DE22 1GB, United Kingdom.
56 57	24	Tel: +44 (0)7774014822.
58 59 60	25	Email: M.Mina@derby.ac.uk

26 ABSTRACT

Studies examining acute, high-speed movement performance enhancement following intense muscular contractions (frequently called 'post-activation potentiation'; PAP) often impose a limited warm-up, compromizing external validity. In the present study the effects on countermovement vertical jump (CMJ) performance of back squat exercises performed with or without elastic bands during warm-up were compared. After familiarization, fifteen active men visited the laboratory on two occasions under randomized, counterbalanced experimental squat warm-up conditions: (1) free-weight resistance (FWR) and (2) variable resistance (VR). After completing a comprehensive task-specific warm-up, three maximal CMJs were performed followed by three back squat repetitions completed at 85% of 1-RM using either FWR or VR. Three CMJs were then performed 30 s, 4 min, 8 min and 12 min later. During CMJ trials, hip, knee and ankle joint kinematics, ground reaction force data and vastus medialis, vastus lateralis and gluteus maximus electromyograms (EMG) were recorded simultaneously using 3D motion analysis, force platform, and EMG techniques, respectively. No change in any variable occurred after FWR (p>0.05). Significant increases (p<0.05) were detected at all time points following VR in CMJ height (5.3-6.5%), peak power (4.4-5.9%), rate of force development (12.9-19.1%), peak concentric knee angular velocity (3.1-4.1%) and mean concentric vastus lateralis EMG activity (27.5-33.4%). The lack of effect of the free-weight conditioning contractions suggests that the comprehensive task-specific warm-up routine mitigated any further performance augmentation. However, the improved CMJ performance following the use of elastic bands is indicative that specific alterations in force-time properties of warm-up exercises may further improve performance.

Keywords: elastic bands, PAP, conditioning activity, explosive strength, kinetics, kinematics

INTRODUCTION

Pre-exercise (i.e. warm-up) routines are typically designed to precondition the neuromuscular system to enhance performance and reduce injury risk during subsequent high-intensity physical activities.¹⁻³ Performing maximal or near-maximal muscular contractions during a warm-up routine are important as they can induce short-term increases in force production and physical performance⁴ through a number of mechanisms including, but not limited to, increases in muscle temperature,⁵ reductions in muscle thixotropy or viscosity,³ increases in myofilament calcium sensitivity,⁶ an increased neural drive (leading to higher-frequency motor unit discharge) and optimization of motor control strategies.⁷ Such changes lead to an increased mechanical power output (i.e. above previous maximal voluntary capacity), a state often referred to as post-activation potentiation (PAP) but which may not be synonymous with 'classic' PAP, which refers to an increase in muscular force production during an electrically elicited (twitch) contraction.⁸ Regardless of the mechanism, short-term improvements in performance (i.e. post-activation performance enhancements [PAPE])⁹ are commonly reported following intense muscular contractions that have important implications for the design of warm-up strategies.

The acute augmentation of physical performance has been explored using different warm-up strategies including light muscle stretching, cycling, running and sub-maximal repetitions of the primary task¹⁰ or no warm-up at all.⁸ Consequently, a "comprehensive task-specific" warm-up (including progressively intense task-specific conditioning activities) is often not provided prior to the specific activity being tested. Although warm-up strategies adopted to potentiate muscular force production have been shown to enhance athletic performance following a conditioning activity, it is unclear whether the enhancement of athletic performance observed is a consequence of acute neuromuscular alteration relating to the conditioning
 activity, or whether it simply reflects a standard warm-up itself.¹¹

Heavy resistance exercise has been shown to acutely potentiate muscle force output, at least when a comprehensive task-specific warm-up is not completed,^{2,8,12} however force production can also be reduced as a result of fatigue or coordination interference (i.e. perseveration) processes, which may mask any potentiating effects.¹ Some studies have reported that vertical jump performance enhancements can be detected after only 20 s¹³ and 90 s¹⁴ following maximal isometric squats and heavy box squats, respectively. Findings from these studies are indicative that effects may be detected within the time course of "classic" PAP observed using muscle twitch examinations.⁸ Nonetheless, a meta-analysis of the literature revealed that minimal performance enhancement was likely when the rest period was less than 2 min, whereas longer rest periods of 3-7 min were more beneficial.¹⁵ The equivocal findings likely result from disparate study methodologies including types of conditioning activity (i.e. movement-pattern specificity), performance tasks, delay between the conditioning activity and performance testing, study participant characteristics (e.g. experienced/novice lifters) and warm-up performed, which limit our understanding of the potentiating effects of these warm-up strategies.

The countermovement vertical jump (CMJ) task is commonly performed in sport but is also a model commonly used to test power and muscle function in clinical research environment. Various high-intensity exercise types have been performed before maximal CMJ tests including resistance-, plyometric-, and electrical muscle stimulation-based exercises.^{16,18} The back squat exercise is a fundamental exercise for the development of lower-limb strength and power¹² and its use during a warm-up has been reported to improve subsequent functional

performance including CMJ height;^{1,4} this enhancement is commonly attributed to the PAP effect. However, maximal voluntary muscle activity occurs only during a short period in the early ascending (concentric) phase, near the "sticking point" in successful maximal (1-RM) back squat attempts. The larger internal and smaller external moment arms developed at the hip and knee joints (resulting in a greater mechanical advantage) combined with the optimized force-length characteristics of lower-limb muscles, ensures that only a submaximal muscle activation is needed for successful completion of the remaining part of the lift.¹⁸ Thus, theoretically, variations of the exercise that evoke a greater muscle activation throughout the lift could result in a greater warm-up (i.e. PAPE) effect and improve CMJ performance. A possible means to alter the loading characteristics of the squat lift is the use of elastic bands to reduce the external load in the deepest part of the squat while increasing external load when the joints are more extended, the internal moment arms are greater and optimal muscle lengths are achieved.^{18,19} Previous studies comparing elastic bands to free-weight squats for muscle activities (EMG), kinematics and kinetics has shown significantly higher EMG, movement velocity, and external power in the first guarter of the eccentric phase and the last guarter of the concentric phase of the squat exercise when using elastic bands.¹⁹ Accordingly, it has been found that preconditioning contractions using elastic bands significantly increased subsequent 1-RM squat test performance without detectable changes in knee extensor muscle activity or knee flexion angle, although eccentric and concentric velocities were reduced.²

Accordingly, elastic bands can be used to increase resistance in ranges of motion where the muscles can produce the greatest relative force as well as unload the system where muscle forces are compromized, and thus allow a larger overall impulse to be produced. Given the possibility for higher muscle activation and greater total work done during the lift, it might be hypothesized that these conditions would allow for a greater potentiating effect.

Individuals incorporating the use of elastic band-based strategies into a warm-up routine may observe an acute enhancement of performance, and thus benefit from a greater mechanical stimulus during training.² However, a common limitation in the literature is that minimal or no warm-up has been provided before imposing the conditioning activity⁸ limiting the comprehensive applicability and external validity of the data. Therefore, the purpose of this study was to compare the influence of free-weight resistance (FWR) and variable resistance (VR; imposed through elastic bands) squat exercises following a task-specific comprehensive warm-up on subsequent CMJ performance at different post-conditioning time points (i.e. 30 s, 4 min, 8 min, and 12 min). It was hypothesized that (i) FWR and VR would enhance subsequent CMJ performance; however the variation in resistance imposed by the elastic bands during the squat lift would (ii) further enhance subsequent CMJ performance, (iii) alter CMJ kinetic and kinematic parameters (i.e. peak power, peak eccentric kinetic energy, impulse- and time-based descent-to-ascent asymmetry indexes, vertical stiffness (K_{vert}), rate of force development (RFD), hip, knee and ankle joint kinematics), and (iv) increase the muscle activity of the lower-limb extensor muscles more than squatting without elastic bands.

0 141

142 MATERIALS AND METHODS

Participants

Fifteen active men (age = 21.7 ± 1.1 y, height = 1.8 ± 1.9 m, mass = 77.6 ± 2.6 kg) with ≥ 5 y experience with heavy weight training of varying levels (from regional to elite) and training backgrounds volunteered to participate after providing written informed consent and completing a pre-test medical questionnaire. The participants' training protocols involved resistance training, sprint running, power exercises, dynamic/explosive exercises, agility drills and other specific exercises relevant to their sports. The participants had no recent illness or

lower-limb injury, were instructed to maintain normal eating and drinking habits throughout the study, and avoided strenuous exercise and stimulant use at least 48 h prior to testing. Ethical approval was granted by the ethics committee at the University of Thessaly, Greece, with the study conducted in accordance with the Declaration of Helsinki. Effect size (ES) values (Cohen's d) were calculated from mean changes in variables (jump height, power, RFD and EMG) from previous studies using similar methods. To ensure an adequate population to reach statistical power (set at 0.8) was recruited, effect sizes were initially calculated from related research²¹⁻²³ for jump height (ES = 1.48), power (ES = 1.0), RFD (ES = 1.29), and EMG (ES = 1.2). To ensure an adequate sample, the measure with the smallest ES (power, 1.0) was used to calculate sample size. The analysis revealed that the initial sample size required for statistical power was 14; thus, considering the possibility of participant withdrawal and data loss, 18 participants were recruited with 15 participants completing the study.

Protocol

Overview

A randomized, cross-over design was implemented to compare CMJ performance following two warm-up conditions: free-weight resistance (FWR) or variable resistance (VR) back squat Participants completed a familiarization session one week prior to the two exercise. experimental sessions, each separated by 72 h and performed at the same time of the day. During familiarization, anthropometric characteristics were recorded, one-repetition maximum (1-RM) back squat load was determined, and the participants were familiarized with all experimental procedures. During experimental conditions, following the comprehensive task-specific warm-up (described later), the participants performed three pre-intervention CMJs followed by back squats at 85% of 1-RM using either FWR or VR warm-up. CMJ trials were then performed at 30 s, 4 min, 8 min and 12 min after the intervention. Peak power output,

peak eccentric kinetic energy, impulse- and time-based descent-to-ascent asymmetry indexes,
peak normalized (to body weight) K_{vert} and RFD, peak knee flexion angle, peak eccentric and
concentric knee angular velocities, peak and mean eccentric and concentric electromyograms
(EMG; vastus lateralis (VL), vastus medialis (VM), gluteus maximus (Glut)), and jump height
were measured during all CMJ trials (described later).

181 Familiarization session and one-repetition (1-RM) squat lift test

The 1-RM back squat protocol was adopted from Sheppard and Triplett et al.²⁴ Participants initially performed a 5 min cycling warm-up (Monark 874E, Varberg, Sweden) at 65 rpm with a 1-kg resistance load followed 2 min later by 2 sets of 10 back squat repetitions using an unloaded 20-kg Olympic bar. The participants then completed 8-10 repetitions of the squat lift exercise at 50% of their estimated 1-RM load before the load was increased by 20% for 3-5 repetitions, and by a further 20% for 2-3 repetitions with a 2 min rest between sets. The load was finally increased by 5% movements with 2-4 min rest between lifts until participants failed to complete the lift; the previous successful attempt was recorded as their 1-RM load. To ensure correct technique, participants were instructed to place the bar above the posterior deltoids at the base of the neck and position the feet shoulder width apart with the toes pointed slightly outward and attempt to squat to a position where the knee was flexed to $\sim 90^{\circ}$ before returning to a standing position. This was visually assessed by an experienced, certified British Amateur Weight Lifting Association (BAWLA) spotter throughout all testing procedures to ensure correct technique and safety during the lifts, with participants receiving strong verbal encouragement to promote maximal effort.

Comprehensive task-specific warm-up and countermovement jump trials

During the experimental trials the participants performed a comprehensive task-specific warm-up consisting of 5 min of cycling followed by five continuous unloaded squats (i.e. non-jumping) at a rhythm of 2 s/2 s (eccentric/concentric) and a further 5 squats at a rhythm of 1 s/1 s after a 30 s rest. After 20-s rest, five continuous CMJs were performed at ~70% of the participants' perceived maximum and, after a further 30 s rest, maximal CMJs were performed every 30 s until three consecutive jumps were within 3% of jump height (4-7 jumps were performed in all trials). The CMJ was performed from a stationary upright standing position with hands positioned on the hips, making a preliminary downward movement with the hips and knees flexed, and immediately jumping vertically up as high possible.¹² Two minutes after the completion of the warm-up, three maximal pre-intervention CMJ trials were performed to establish baseline (i.e. after warm-up) performance. A conditioning set of three repetitions of back squats at 85% of the previously determined 1-RM using either FWR or VR (described later) was then performed before the participants completed three CMJs 30 s, 4 min, 8 min and 12 min (see Table 1) later with participants receiving verbal encouragement to jump as high as possible. The post-intervention intervals were selected from previous data describing the time-course of the performance augmentation (PAP) response.^{21,25} Table 1 about here Conditioning activities: free-weight resistance (FWR) and variable resistance (VR) During the FWR condition, the load was adjusted to 85% of the previously determined 1-RM load with the participants performing one set of three-repetition back squats. In the VR warm-up condition, 35% of the total load was generated from band resistance. To ensure a similar load of 85% 1-RM across FWR and VR conditions, mechanical properties of the bands were

determined to allow the band resistance to generate 35% of the total load. Half of the band's resistance was subtracted from the total free-weight load to ensure the elastic bands did not have substantially different average resistance compared with the FWR condition, thus both the FWR and VR warm-up conditions were equalized, as previously reported.^{20,22} The participants stood on a force platform with 85% 1-RM load to determine the combined load (kg), the bar was then unloaded to adjust the band tension. The elastic bands were anchored to the floor with custom-made weight stands and attached equidistant to the ends of the Olympic bar to ensure the participant's stability. The thickness and lengths of the elastic bands were adjusted so that: (i) the tension in the bands increased the ground reaction force (measured by force platform) by 35% of the 85% load when the participants were standing, but (ii) bands were slack in a full squatting position and thus provided no additional loading. The linear force-length properties of the bands ensured, therefore, that the average load during the lift equated to 35% of the total load. For example, a 100-kg load in the FWR condition would require 35-kg (35%) to be generated from the bands. Half of the 35-kg load (i.e. 17.5 kg) was removed from the bar with the 35-kg resistance added from the bands providing a total load of 117.5 kg in the standing position. As band tension reduced as the participant squats, 35 kg of load was removed leaving the 82.5 kg from the bar in the full squatted position. Thus, the average loading throughout the lift in this example is 100 kg, identical to the FWR condition whilst enabling 35% to be generated by band tension.

47 243

 Kinetic and kinematic analyses

Kinematic data were collected during the CMJs using a Vicon motion analysis system (TSeries, Oxford Metrics LTDA, Oxford, UK) with 10 cameras operating at 100 Hz surrounding
two force platforms (Bertec, FP4060-10-2000, Bertec Corporation, Columbus, OH, USA).
Ground reaction forces were sampled at 1000 Hz and time-synchronized with the Vicon system

(see Figure 1). The data were then filtered using Woltring's quantic spline algorithm²⁶ with a mean squared error setting of 15 before running the Plug-In-Gait biomechanical model (Vicon Plug-in-Gait, Oxford Metrics). The procedures identified by Davis et al.²⁷ were followed to define Cardan angles and to reconstruct a system of embedded coordinates from the marker set to 0° at the three joints of the lower extremities (hip, knee and ankle) in a standing position. Lower-limb kinetic and kinematic data were captured by placing 16 reflective markers over the pelvis, left and right thigh, left and right shank in a straight line, and the left and right foot at a right angle to the leg. Data were analyzed using Vicon Nexus (v.2.3) software to determine peak hip, knee and ankle flexion angle and angular velocity data during the pre- and postintervention CMJ trials (see Figure 1).

Figure 1 about here

All jumps were performed from the standing position with each foot in parallel on two force platforms providing a separate vet time-synchronized measurement of the force data for each leg. The participant's body weight was calculated by averaging the vertical force from each platform when the participants were stationary. The initiation of the jump (i.e. the beginning of the eccentric phase) was identified as the point when the ground reaction force (N) decreased 2 standard deviations (SD) below the mean baseline force. The vertical ground reaction force was integrated using the trapezoid method during the eccentric and concentric phases of the jump. The net impulse was calculated independently and summed from the left and right force platforms. Ground reaction forces were directly quantified by integrating the applied force over time (i.e. impulse), which is equivalent to the change in momentum of the body:

273 where J = impulse, F = force, t = time and $\Delta p =$ change in momentum.

 $J = \int F dt = \Delta p$

2	
3 4	274
5 6	275
7 8 0	276
9 10 11	277
12 13	278
14 15	279
16 17	280
18 19 20	281
20 21 22	282
23 24	283
25 26	284
27 28 20	285
29 30 31	205
32 33	200
34 35	287
36 37	288
38 39	289
40 41 42	290
42 43 44	291
45 46	292
47 48	293
49 50	294
51 52	295
55 55	296
56 57	297
58 59	298

1

The take-off velocity was determined from impulse by dividing by body mass, and the jump 275 height was calculated using standard equations for motion.²⁸ To calculate power, the impulse-276 momentum approach was used. Since the force, mass and initial velocity conditions were 277 known, instantaneous velocity could be calculated. The instantaneous power was calculated 278 as force \times velocity and the peak values were determined for the propulsive phase of the CMJ: 279 $V_{(0)} = 0$ 280 $F(i)t = m(v_{(i+1)} - v_{(i)})$ 281 $\Delta v = (F_{(i)}t)/m$ 282 $P_{(i)} = F_{(i)} \times V_{(i)}$ 283 where F =force, t = 1/sampling frequency, m = mass of body, load, v = velocity, and P =284 power. 285 286 The peak eccentric kinetic energy (KE) developed during the jumps was calculated as: 287 $KE = \frac{1}{2}mv^2$ 288 where m is the participant's mass and v is the velocity of the countermovement phase. 289 290 The impulse-based asymmetry index was calculated by dividing the negative and positive 291 impulses, where the negative impulse describes the impulse that negatively accelerates the 292 body downwards and the positive impulse accelerates the body upwards. The index was 293 calculated to estimate the efficiency of the metabolic energy conversion into mechanical work 294 (i.e. storage of elastic energy during eccentric contraction) performed during the CMJ from the 295 force applied by the body to the ground²⁹ and subsequently released energy during the 296

298 of times A + B, where A is the time from force first rising above 1 body weight to the peak

concentric phase of the SSC. The time-based asymmetry index was calculated as the quotient

2		
3 4	299	vertical force and B is the time from peak force until force drops below 1 body weight. K _{vert}
5 6	300	was calculated by dividing the peak vertical ground reaction force by the maximal vertical
7 8	301	displacement of the center of mass during contact with the ground ³⁰
9 10 11	302	$K_{vert} = F_{max} / \Delta y$
12 13	303	where F_{max} = maximum vertical force, and Δy = maximum vertical displacement of the center
14 15	304	of mass. The vertical displacement was determined by the double integration of the vertical
16 17 19	305	force trace according to methods of Cavagna. ³¹
19 20	306	
21 22	307	The peak RFD (normalized to body weight) was calculated from the initiation of the jump (i.e.
23 24	308	first rise in force during the eccentric phase) using the average force-time curve with a 50-ms
25 26 27	309	time window.
28 29	310	
30 31	311	Muscle activity (electromyography; EMG)
32 33 34	312	EMG data were collected wirelessly using a Myon MA-320 EMG system (Myon AG,
35 36	313	Schwarzenberg, Switzerland) from vastus lateralis (VL), vastus medialis (VM) and gluteus
37 38	314	maximus (Glut). The skin was shaved, abraded and cleansed with alcohol before bipolar
39 40	315	adhesive surface electrodes (Noraxon Dual Electrodes, Ag-AgCl, Noraxon USA, Inc,
41 42 43	316	Scottsdale, AZ) were placed over the muscle belly with an inter-electrode distance of 2 cm
44 45	317	according to SENIAM guidelines. EMG data were sampled at 2000 Hz and imported into
46 47	318	ProEMG software (version 4.1) and filtered using a Butterworth (20-500 Hz bandpass) filter
48 49 50	319	before using a symmetric moving root-mean-square algorithm with a 50-ms sampling window.
50 51 52	320	The Myon EMG software was integrated with an optimal tracking device for synchronization
53 54	321	between the systems (Vicon motion analysis system, Oxford, UK). The normalized EMG
55 56 57	322	amplitude during isometric squat lifts (% maximal voluntary contraction [MVC]) for each
58 59		

muscle was used as a measure of neuromuscular activity during the jumps (see Figure 1), with

peak and mean EMG activity recorded during the eccentric and concentric phases.

Data analyses

All data were analyzed using SPSS statistical software (version 24.0; IBM, Chicago, IL, USA); all data are presented as mean \pm SE. Normal distribution was assessed using Shapiro-Wilks test; no significant difference (p > 0.05) was detected in any variable indicating that all data sets were normally distributed. Separate multivariate analyses of variance (MANOVAs) were used to compare (a) jump height and peak power, and (b) EMG. Where significant differences were detected, separate two-way repeated measures ANOVAs (time \times condition) were used to determine differences in (a) jump height, (b) peak power, (c) peak eccentric kinetic energy, (d) impulse- and time-based descent-to-ascent asymmetry indexes, (e) peak normalized RFD, (f) peak hip, knee and ankle flexion angle, (g) peak eccentric and concentric hip, knee and ankle angular velocities, (h) peak and mean eccentric and concentric EMG activities during CMJ trials. Significance was accepted at p < 0.05 for all tests.

Reliability

Reliability for all measures was determined during the pre-intervention vertical jumps from the VR and FWR warm-up conditions. No significant differences (p > 0.05) were detected in any measure and high intraclass correlation coefficients (ICCs) calculated for jump height (0.95), peak power (0.98), peak eccentric kinetic energy (0.99), impulse- (0.96) and time-based (0.91)asymmetry indexes, K_{vert} (0.81), peak RFD to 50 ms (0.92), peak hip, knee and ankle flexion angle ranged from 0.67 to 0.96, peak angular velocities ranged from 0.76, 0.95, 0.85 to 0.85, 0.95 0.79 for hip, knee and ankle, respectively. ICCs for the EMG data ranged from 0.73 to 0.89, 0.85 to 0.92, 0.85 to 0.92 for VL, VM and Glut, respectively. Coefficients of variation

(CoV) expressed as a percentage of the mean were also calculated for jump height (8.0%), peak power (6.2%), peak eccentric kinetic energy (8.5%), impulse- (4.9%), and time-based (14.6%) asymmetry indexes, K_{vert} (8.7%), peak RFD to 50 ms (12.5%), peak hip, knee and ankle flexion angle ranged from 3.8% to 7.6%, peak angular velocities ranged from 5.6%, 5.2%, 14.8% to 5.0%, 3.4%, 6.3% for hip, knee and ankle, respectively. CoVs for EMG data 9.0% to 14.3%, 11.3% to 14.1%, 14.9% to 22% for VL, VM and Glut, respectively.

RESULTS

In the FWR condition, no significant changes (p > 0.05) were found in jump height (range = 3.0 ± 2.0 % to 4.9 ± 2.2 %) at any time point compared with pre-intervention data (see Figure 1). Also, no significant changes ($p \ge 0.05$) were observed in peak power ($0.1 \pm 2.4\%$ to $3.6 \pm$ 1.6%), peak eccentric kinetic energy $(0.5 \pm 4.6\% \text{ to } 4.9 \pm 3.6\%)$, impulse- $(0.6 \pm 1.6\% \text{ to } 2.0 \pm 1.6\%)$ 1.9%) and time-based $(4.5 \pm 7.0\% \text{ to } 14.8 \pm 8.4\%)$ asymmetry indexes, $K_{vert} (3.1 \pm 4.3\% \text{ to } 5.8\%)$ $\pm 4.1\%$) or peak normalized RFD (3.1 $\pm 6.1\%$ to 11.8 $\pm 8.4\%$) at any time point (see Table 2). No changes (p > 0.05) were detected in peak eccentric hip $(0.5 \pm 1.7\% \text{ to } 2.6 \pm 1.9\%)$, knee $(0.5 \pm 2.1\%$ to $2.6 \pm 2.2\%)$, ankle $(2.2 \pm 5.4\%$ to $9.0 \pm 5.0\%)$ or concentric hip $(1.2 \pm 2.1\%$ to $3.7 \pm 2.0\%$), knee (0.5 ± 1.7% to 1.7 ± 2.0%), ankle (1.4 ± 2.1% to 4.7 ± 2.6%) angular velocities, or peak hip $(1.5 \pm 1.3^{\circ} \text{ to } 3.4 \pm 1.2^{\circ})$, knee $(0.1 \pm 1.2^{\circ} \text{ to } 1.7 \pm 1.9^{\circ})$, ankle $(0.1 \pm 1.2^{\circ})$ 0.6° to $0.6 \pm 0.5^{\circ}$) flexion angle (see Table 3). Furthermore, no changes in peak or mean eccentric EMG eccentric activity (p > 0.05) in VL (peak = $2.4 \pm 3.8\%$ to $7.2 \pm 5.2\%$; mean = $0.7 \pm 6.4\%$ to $7.3 \pm 5.7\%$), VM (peak = $0.6 \pm 3.2\%$ to $8.3 \pm 5.0\%$; mean = $8.9 \pm 4.5\%$ to 10.9 \pm 3.2%), Glut (peak = 0.9 \pm 6.0% to 8.7 \pm 3.8%; mean = 2.3 \pm 6.0% to 10.7 \pm 7.4%) or concentric EMG in VL (peak = $0.4 \pm 10.3\%$ to $9.4 \pm 8.3\%$; mean = $2.2 \pm 8.6\%$ to $7.0 \pm 6.9\%$), VM (peak = $0.5 \pm 4.6\%$ to $7.1 \pm 5.2\%$; mean = $1.2 \pm 7.1\%$ to $9.5 \pm 5.4\%$) or Glut (peak = 1.3 $\pm 5.7\%$ to $10.4 \pm 5.6\%$; mean = $2.1 \pm 6.6\%$ to $8.3 \pm 8.9\%$) were detected (see Table 4).

1		
2 3	272	
4	3/3	
5 6 7	374	Table 2 about here
7 8 9	375	Figure 2 about here
10 11	376	
12 13 14	377	In the VR condition, significant increases ($p < 0.05$) in CMJ height were detected at 30 s (5.9
14 15 16	378	\pm 1.2%), 4 min (5.6 \pm 1.8%), 8 min (6.5 \pm 2.6%) and 12 min (5.3 \pm 2.5%) time points compared
17 18	379	with pre-intervention data (see Figure 2). Significant increases ($p < 0.05$) were also observed
19 20 21	380	in peak power at 30 s (4.7 \pm 1.2%), 4 min (5.9 \pm 1.3%), 8 min (4.4 \pm 1.7%) and 12 min (4.8 \pm
21 22 23	381	1.7%) time points compared to pre-intervention data. These changes in CMJ height and power
24 25	382	were also statistically different to FWR ($p < 0.05$). Similarly, significant increases ($p < 0.05$)
26 27	383	were found in peak normalized RFD at 30 s ($18.9 \pm 7.8\%$), 4 min ($12.9 \pm 5.9\%$), 8 min ($19.1 \pm$
28 29 30	384	5.0%) and 12 min (16.0 \pm 8.1%) compared to pre-intervention data. However, no significant
31 32	385	change ($p > 0.05$) in peak eccentric kinetic energy ($0.4 \pm 4.8\%$ to $5.2 \pm 4.8\%$) or impulse- (1.4
33 34	386	\pm 1.5% to 4.6 \pm 2.4%) or time-based (7.4 \pm 11.7% to 13.0 \pm 12.5%) asymmetry indexes, K_{vert}
35 36 37	387	$(6.6 \pm 4.5\%$ to $8.9 \pm 3.7\%)$ were found following the VR warm-up condition at any time point
38 39	388	(see Table 2).
40 41	389	
42 43	390	Table 3 about here
44 45 46	391	
47 48	392	No significant change in peak hip $(1.3 \pm 1.3^{\circ} \text{ to } 1.9 \pm 1.5^{\circ})$, knee $(0.9 \pm 2.9^{\circ} \text{ to } 4.1 \pm 3.0^{\circ})$,
49 50	393	ankle $(0.9 \pm 0.4^{\circ} \text{ to } 1.4 \pm 0.7^{\circ})$ flexion angles were observed in VR at any time point. Similarly,
51 52 53	394	no changes ($p > 0.05$) were found at any time point (see Table 3) in peak eccentric hip (0.2 ±
54 55	395	2.2% to 2.5 \pm 2.2%), knee (0.04 \pm 1.7% to 2.6 \pm 2.5%), ankle (0.1 \pm 6.6% to 6.7 \pm 8.7%) or
56 57	396	concentric hip $(1.5 \pm 2.4\%$ to $3.6 \pm 2.1\%)$ or ankle $(1.1 \pm 2.0\%$ to $3.5 \pm 2.1\%)$ angular velocities
58 59 60	397	or peak or mean eccentric EMG amplitudes for VL (peak = $0.5 \pm 4.3\%$ to $3.1 \pm 4.3\%$, mean =

398	$4.9 \pm 6.0\%$ to $9.2 \pm 6.5\%$), VM (peak = $2.1 \pm 4.6\%$ to $9.6 \pm 4.0\%$, mean = $4.9 \pm 5.4\%$ to $6.7 \pm 6.0\%$
399	5.8%) or Glut (peak = $2.2 \pm 6.3\%$ to $4.6 \pm 6.3\%$, mean = $3.5 \pm 7.4\%$ to $4.9 \pm 6.5\%$). However,
400	a significant increase ($p < 0.05$) was found in peak concentric knee angular velocity at 30 s (3.1
401	$\pm 1.4\%$), 4 min (4.1 $\pm 1.7\%$), 8 min (3.2 $\pm 1.0\%$) and 12 min (3.1 $\pm 1.5\%$) and mean concentric
402	VL EMG activity at 30 s (28.1 \pm 10.5%), 4 min (31.5 \pm 11.0%), 8 min (33.4 \pm 15.9%) and 12
403	min (27.5 \pm 14.5%) compared to pre-intervention data. No changes ($p > 0.05$) in mean
404	concentric VM ($3.7 \pm 8.0\%$ to $12.7 \pm 8.6\%$) or Glut ($0.3 \pm 10.4\%$ to $7.0 \pm 7.5\%$) EMG or peak
405	concentric VL ($0.6 \pm 5.8\%$ to $4.5 \pm 4.7\%$), VM ($0.3 \pm 5.1\%$ to $9.2 \pm 4.1\%$) or Glut ($0.2 \pm 9.2\%$
406	to $7.1 \pm 7.7\%$) EMG were observed at any time point (see Table 4).
407	

Table 4 about here

Significant (p < 0.05) correlations were observed between the change in CMJ height (preintervention to 8 min post-intervention, i.e. where the greatest mean increase in jump height occurred) and changes in peak power (r = 0.82) during VR. No significant correlations (p >0.05) were found between change in CMJ height and changes in peak normalized RFD (r =0.27), peak knee angular velocity (r = -0.21), mean concentric VL EMG (r = 0.17) or peak eccentric kinetic energy (r = 0.32).

DISCUSSION

418 The primary aim of the present study was to assess the magnitude and time-course of changes
51 in countermovement vertical jump (CMJ) performance after traditional free-weight (FWR) and
53 variable (VR) resistance squat exercises were performed following a comprehensive task54 variable (VR) resistance squat exercises were performed following a comprehensive task55 specific, warm-up routine. The first hypothesis can be partially accepted as the lack of change
58 in any measure following the FWR condition suggests that no additional benefit (i.e.

PAP/PAPE effect) was derived from the inclusion of intense loading from FWR exercise (i.e. the conditioning activity), contrary to the improvement in jump height following the use of elastic bands. This finding contrasts those of previous studies where the performance of heavy squat lifts increased CMJ height,^{21,32} and other literature reporting an increase in tasks including sprint running performance.³³ However, those previous studies either did not report the use of other warm-up activities or only included a light cardiovascular warm-up rather than a more comprehensive task-specific warm-up including progressively intense task-specific muscular contractions. The current finding of a lack of effect of a back squat conditioning activity after a comprehensive task-specific warm-up (see Figure 2) is, however, consistent with a previous report of an absence of change in vertical jump performance when dynamic warm-up exercises were employed prior to a set of back squats.¹⁷ These data are indicative that a lack of a comprehensive task-specific warm-up may enable further augmentation of performance after squats were performed, but may be of limited relevance to athletes, strength trainers and recreational exercisers who would customarily perform a thorough warm-up. That is, the high-intensity conditioning activities might only increase performance when the warm-up would otherwise be insufficient to promote maximal performance. Collectively, these findings indicate that the previously reported 'potentiating' effects of heavy free-weight back squat exercise on subsequent CMJ performance^{16,21} may be a consequence of study design, where the limited use of warm-up protocols provided an opportunity for further performance augmentation after the baseline tests. Furthermore, inconsistencies in PAP responses^{12,16,17} may depend on fatigue-potentiation or perseveration-potentiation interactions and their influence on subsequent performance therefore new strategies for designing warm-up protocols and optimal recovery periods following conditioning contractions are vital in order to induce a potentiation effect.

Despite FWR squat lifts having no effect on CMJ performance, a significant increase in jump height was achieved following the VR conditioning activity at all time points (30 s, 4 min, 8 min and 12 min; see Figure 2), which suggests a prolonged 'potentiating' effect was evoked, i.e. post-activation performance enhancement; PAPE.⁹ Thus, the second experimental hypothesis, that jump height would be increased following the VR intervention, can be accepted. These data are consistent with a previous study³⁴ in which box squats incorporating elastic band resistance acutely increased power output during subsequent CMJ tasks. However, in the present study it was shown that this effect can be evoked even after completion of a comprehensive task-specific warm-up, which was not included in previous studies. Although each maximal CMJ may possibly potentiate the next one, no significant improvement occurred in the FWR condition, therefore these jumps were unlikely to explain the increased performance in VR. Previous studies showed that only seconds or a few minutes are needed to recover from a short bout of maximal-effort exercise (e.g. less than 1 min for recovery from a maximal squat³⁵ or bench press lifts³⁶), thus it is unlikely that fatigue is a factor influencing the findings of the present study as a significant increase was observed across all time points. The use of elastic bands reduces the effective load near the "sticking point" in the early concentric phase of the squat lift but then allows for greater loading later in the lift as the effective mechanical advantage is increased.² The ability for muscles to operate closer to their maximum force capacity through a greater proportion of the lift may therefore enhance subsequent muscle force output and elicit a greater dynamic muscle performance (i.e. increase in CMJ height), even when a comprehensive task-specific warm-up is already completed. Collectively, these data indicate that the use of elastic bands, which alter the loading strategy during the lift, provides a more effective warm-up than either warm-up alone or warm-up that also includes traditional free-weight resistance exercises.

In the present study significant changes in force production (peak power and RFD; see Table 2) at all time points in the VR warm-up condition were consistent with the changes in jump height. However, peak hip, knee and ankle, flexion angle, peak eccentric kinetic energy, the impulse- and time-based asymmetry indexes remained unchanged and no change was detected in K_{vert} (see Tables 2 and 3), which is consistent with previous research.³⁰ Accordingly, changes in jump kinematics cannot explain the changes in force production or jump height. The third hypothesis, that both kinetic and kinematic parameters would be altered by elastic band-resisted squat lifts, can therefore only be partially accepted. The changes in peak jump power were significantly correlated with the changes in CMJ height, however a poor relationship was identified between changes in RFD and CMJ height. This latter finding is consistent with a previous report³⁷ in which stretch-shortening cycle test performances were not statistically related to RFD measured during the test. The poor relationship may be partly explained by the participants being well strength-trained yet relatively untrained in explosive power-based exercises, and thus unable to rapidly reach peak force.³⁸ However, further research on power-trained athletes is needed to fully elucidate the importance of training status.

A number of mechanisms relating to stretch-shorten cycle efficiency may have contributed to the increased jump height, including a more rapid muscle stretch resulting from force potentiation,³⁹³⁹ greater elastic energy storage in the muscle,⁴⁰ an increased time of muscle activation,^{40,41} an augmented pre-load effect,⁴² force and stiffness augmentation from stretch reflexes,⁴¹ or changes in relative contributions of muscle and tendon allowing the muscle to operate at lower shortening speeds and over shorter distances.⁴³ Whilst it is difficult to assess the effects of each, the peak eccentric kinetic energy and both impulse- and time-based asymmetry indexes remained unchanged after VR, indicating that the total energy available for storage in elastic structures (eccentric kinetic energy), the kinematic pattern adopted to make

use of it (asymmetry indexes),²⁹ and the time for force application and likely contribution of stretch reflexes, were also unchanged. Nonetheless, increases in peak power and concentric knee angular velocity were observed.

Nonetheless, a more plausible explanation for the increase in force production, and thus jump height, may be found in the increased knee extensor muscle activity detected in the concentric phase (VL EMG increased 27.5-33.4% across time points; see Table 4). Thus, the fourth hypothesis, that extensor muscle activity would be increased, can be accepted. The greater increase in EMG activity in VL than VM or Glut is consistent with previous reports of greater VL EMG in the concentric phase of a CMJ after both low- and moderate-intensity squat warm-ups³² and would likely have resulted from an increased motor unit firing frequency.⁴⁴ In fact, Nikolaidou et al.⁴⁵ found that a greater jump height was achieved during CMJ compared to squat jump which was consistent with an increased VL activation during the push off phase. Increased phosphorylation of the myosin light chain leading to an increase in myofilament Ca²⁺ sensitivity and force output (i.e. classic PAP) may have contributed to the increase in CMJ, although it resolves completely within about 5 min¹¹ thus its effect at 4 - 12 min would have been negligible. Although other mechanisms such as increases in muscle temperature⁵ (not examined in the present study) may have contributed to the increase in jump height it remains likely that the change in muscle activation was the major factor influencing the improvement in CMJ performance. The increased muscle activity and consequent increase in peak power output in the concentric phase would have allowed a greater jump height without changes in kinematics or stretch-shorten cycle efficiency (i.e. without changes in eccentric knee angular velocity, eccentric kinetic energy, impulse- and time-based asymmetry indexes or K_{vert}). The most likely explanation for the finding is that the variation in muscle force requirements imposed by the use of elastic band resistance influenced muscle recruitment patterns and

ultimately increased concentric force output.¹⁹ The current findings hint at the possibility that manipulation of loading strategies during warm-up exercises might beneficially alter muscle recruitment amplitude or timing and result in greater performances than achieved through traditional high-intensity, task-specific warm-ups alone. This hypothesis should be explicitly examined in future studies. It is important to note that it was not possible to measure muscle temperatures in the current study. However, muscle temperature would likely have increased substantially during the comprehensive task-specific warm-up so temperature may have remained constant (i.e. in an optimum zone) for a longer time, and any further small increase in temperature from the conditioning activities would have been similar between conditions. This may have allowed the improved muscle activation to result in a greater jump performance and for the increased activation to persist for a longer time. Thus, although it remains to be explicitly examined in future, it can be considered unlikely that muscle temperature differences could explain the between-condition differences in jump performance.

PERSPECTIVE

The completion of brief, high-load free-weight squat exercise following a comprehensive task-specific warm-up failed to alter CMJ height, force/power production or movement pattern. These findings are suggestive that the previously-observed 'potentiating' effect of squat exercise may be a consequence of limited warm-up. The beneficial effect of a free-weight squat strategy to potentiate the system may therefore, be minimal in athletic populations that typically perform high-intensity, task-specific warm-up routines prior to maximal exercise tasks. However, the use of elastic band resistance during these squats resulted in significant increases in jump height, peak power, peak concentric knee angular velocity and peak RFD, as well as increased VL EMG activity in the concentric (propulsive) phase of the jump, which did not return to baseline after 12 min despite a comprehensive task-specific warm-up being

1 ว		
2 3 4	548	completed. The results suggest that the inclusion of tasks in which force-time parameters differ
5 6 7	549	from the outcome task (CMJ in the current study) might evoke positive acute adaptations in
7 8 9	550	addition to those achieved through warm-up alone. Further research is required to determine
10 11	551	whether similar effects are observed following different warm-up strategies and in different
12 13 14	552	athletic tasks, as well as in other populations.
15 16	553	
17 18	554	ACKNOWLEDGEMENTS
19 20 21	555	No funding was received for this work. No conflicts of interest exist.
21 22 23	556	
24 25	557	
26 27 28	558	
29 30	559	
31 32 33	560	
34 35	561	
36 37	562	
38 39	563	
40 41	564	
42 43 44	565	
45 46	566	
47 48	567	
49 50 51	568	
52 53	569	
54 55	570	
56 57	571	
58 59 60	572	

573	RE	FERENCES
574	1.	Tillin NA, Bishop D. Factors modulating post-activation potentiation and its effect on
575		performance on subsequent explosive activities. Sports Med 2009;39:147-166.
576	2.	Mina MA, Blazevich AJ, Giakas G, Kay AD. The influence of variable resistance loading
577		on subsequent free-weight maximal back squat performance. J Strength Cond Res
578		2014;28:2988–2996.
579	3.	Woods K, Bishop P, Jones E. Warm-up and stretching in the prevention of muscular injury.
580		Sports Med 2007;37:1089–1099.
581	4.	Hodgson M, Docherty D, Robbins D. Post activation potentiation: Underlying physiology
582		and implication for motor performance. Sports Med 2005;35:585-595.
583	5.	Racinais S, Oksa J. Temperature and neuromuscular function. Scand J Med Sci Sports
584		2010;20:11–18.
585	6.	Moore RL, Stull JT. Myosin light chain phosphorylation in fast and slow skeletal muscles
586		in situ. Am J Physiol Cell Physiol 1984;247:462–471.
587	7.	Trimble MH, Harp SS. Postexercise potentiation of the H-reflex in human. Med Sci Sports
588		Exerc 1998;30:933–941.
589	8.	Hamada T, Sale DG, MacDougall JD, Tarnopolsky MA. Postactivation potentiation, fibre
590		type and twitch contraction time in human knee extensor muscles. J Appl Physiol
591		2000;88:2131–2137.
592	9.	Cuenca-Fernandez F, Smith IC, Jordan MJ, et al. Nonlocalized postactivation performance
593		enhancement (PAPE) effects in trained athletes: a pilot study. Appl Physiol Nutr Metab
594		2017;42:1122–1125.
595	10.	Jo E, Judelson DA, Brown LE, Coburn JW, Dabbs NC. Influence of recovery duration
596		after a potentiating stimulus on muscular power in recreationally trained individuals. J
597		Strength Cond Res 2010;24:343–347.

1 2			
3 4	598	11.	MacIntosh BR, Robillard ME, Tomaras EK. Should postactivation potentiation be the goal
5 6	599		of your warm-up? Appl Physiol Nutr Metab 2012;37:546-550.
7 8 0	600	12.	Young WB, Jenner A, Griffiths K. Acute enhancement of power performance from heavy
9 10 11	601		load squats. J Strength Cond Res 1998;12:82-84.
12 13	602	13.	Arabatzi F, Patikas D, Zafeiridis A, et al. The post-activation potentiation effect on squat
14 15	603		jump performance: Age and sex effect. Pediatr Exerc Sci 2014;26:187–194.
16 17 19	604	14.	Baker DG. Increase in jump squat peak external power output when combined with
19 20	605		accommodating resistance box squats during contrasting resistance complex training with
21 22	606		short rest periods. J Aust Strength Cond 2008;6:10-18.
23 24	607	15.	Wilson JM, Duncan NM, Marin PJ, et al. Meta-analysis of post activation potentiation and
25 26 27	608		power: Effect of conditioning activity, volume, gender, rest periods, and training status. J
28 29	609		Strength Cond Res 2013;27:854–859.
30 31	610	16.	Gourgoulis V, Aggeloussis N, Kasimatis P, Mavromatis G, Garas A. Effect of a
32 33 34	611		submaximal half-squats warm-up program on vertical jumping ability. J Strength Cond
35 36	612		Res 2003;17:342–344.
37 38	613	17.	Witmer CA, Davis SE, Moir GL. The acute effects of back squats on vertical jump
39 40	614		performance in men and women. J Sports Sci Med 2010;9:206-213.
41 42 43	615	18.	Elliott BC, Wilson GJ. A biomechanical analysis of the sticking region in the bench press.
44 45	616		Med Sci Sports Exerc 1989;21:450–462.
46 47	617	19.	Israetel MA, McBride JM, Nuzzo JL, Skinner JD, Dayne AM. Kinetic and kinematic
48 49 50	618		difference between squat performed with and without elastic bands. J Strength Cond Res
51 52	619		2010;24:190–194.
53 54	620	20.	Wallace BJ, Winchester JB, McGuigan MR. Effects of elastic bands on force and power
55 56 57	621		characteristics during the back squat exercise. J Strength Cond Res 2006;20:268-272.
58 59			

2			
2 3 4	622	21.	Lowery RP, Duncan NM, Loenneke JP, et al. The effects of potentiating stimuli intensity
5 6	623		under varying rest periods on vertical jump performance and power. J Strength Cond Res
7 8	624		2012;26:3320–3325.
9 10 11	625	22.	Mina MA, Blazevich AJ, Giakas G, Seitz LB, Kay AD. Chain-loaded variable resistance
12 13	626		warm-up improves free-weight maximal back squat performance. Eur J Sport Sci
14 15	627		2016;16:932–939.
16 17 19	628	23.	Golas A, Wilk M, Stastny P, Maszczyk A, Pajerska K, Zajac A. Optimizing half squat
19 20	629		postactivation potential load in squat jump training for eliciting relative maximal power in
21 22	630		ski jumpers. J Strength Cond Res 2017;31:3010–3017.
23 24	631	24.	Sheppard JM, Triplett NT. Program design for resistance training. In: Haff GG, Triplett
25 26 27	632		NT, ed. Essentials of strength training and conditioning. 4th ed. Champaign: Human
28 29	633		Kinetics; 2016. 452-456 p.
30 31	634	25.	Kilduff LP, Bevan HR, Kingsley MIC, et al. Post activation potentiation in professional
32 33 34	635		rugby players: Optimal recovery. J Strength Cond Res 2007;21:1134–1138.
35 36	636	26.	Woltring HJ. A fortan package for generalized cross-validatory spline smoothing and
37 38	637		differentiation. Adv Eng Softw 1986;8:104-113.
39 40	638	27.	Davis III RB, Ounpuu S, Tyburski D, Gage JR. A gait analysis data collection and
41 42 43	639		reduction technique. Hum Mov Sci 1991;10:575–587.
44 45	640	28.	Kibele A. Possibilities and limitations in biomechanical analysis of countermovement
46 47	641		jumps: a methodological study. J Appl Biomech 1998;14:105–117.
48 49 50	642	29.	Cavagna GA. Storage and utilization of elastic energy in skeletal muscle. Exerc Sport Sci
51 52	643		Rev 1977;5:89–129.
53 54	644	30.	Boullosa DA, Abreu L, Beltrame LG, Behm DG. The acute effect of different half squat
55 56 57	645		set configurations on jump potentiation. J Strength Cond Res 2013;27:2059-2066.
58 59	646	31.	Cavagna GA. Force platforms as ergometers. J Appl Physiol 1985;39:174-179.
60			

1

2			
3 4	647	32.	Sotiropoulos K, Smilios I, Christou M, et al. Effect of warm-up on vertical jump
5 6	648		performance and muscle electrical activity using half-squats and moderate intensity. J
7 8	649		Sports Sci Med 2010;9:326-331.
9 10 11	650	33.	Chatzopoulos DE, Michailidis CJ, Giannakos AK, et al. Postactivation potentiation effects
12 13	651		after heavy resistance exercise on running speed. J Strength Cond Res 2007;21:1278-
14 15	652		1281.
16 17	653	34.	Buttifant D, Hrysomallis C. Effect of various practical warm-up protocols on acute lower-
18 19 20	654		body power. J Strength Cond Res 2015;29:656–660.
20 21 22	655	35.	Hitchcock HC. Recovery of short-term power after dynamic exercise. J Appl Physiol
23 24	656		1987;67:677-681.
25 26 27	657	36.	Matuszak ME, Fry AC, Weiss LW, Ireland TR, McKnight MM. Effect of rest interval
27 28 29	658		length on repeated 1 repetition maximum back squat. J Strength Cond Res 2003;17:634-
30 31	659		637.
32 33	660	37	. Wilson G, Lyttle A, Ostrowski K, Murphy A. Assessing dynamic performance: A
34 35 26	661		comparison of rate of force development tests. J Strength Cond Res 1995;9:176–181.
36 37 38	662	38.	Moir G. Sanders R. Button C. Glaister M. The influence of familiarization on the reliability
39 40	663		of force variables measured during unloaded and loaded vertical jumps. I Strength Cond
41 42	664		Res 2005.19.140_145
43 44	665	30	Ressier DE Herzog W Considerations on the history dependence on muscle contraction
45 46 47	003	39.	L Appl Device 2004:06:410, 427
47 48	666		J Appl Physiol 2004,96:419–427.
49 50	667	40.	Finni T, Ikegawa S, Lepola V, Komi PV. Comparison of force-velocity relationships of
51 52	668		vastus lateralis muscle in isokinetic and in stretch-shortening cycle exercises. Acta Physiol
53 54 55	669		Scan 2003;177:483–491.
56 57	670	41.	Bobbert MF, Gerritsen KGM, Litjens MCA, van Soest AJ. Why is countermovement jump
58 59 60	671		height greater than squat jump height? Med Sci Sports Exerc 1996;28:1402-1412.

2		
3 4	672	42. Finni T, Ikegawa S, Komi PV. Concentric force enhancement during human movement.
5 6	673	Acta Physiol Scan 2001;173:369–377.
7 8	674	43. Hill AV. The heat of shortening and the dynamic constants of the muscle. Proc R Soc B
9 10 11	675	Biol Sci 1938;843:136–195.
12 13	676	44. Heckman CJ, Enoka RM. Motor Unit. Compr Physiol 2012;2:2629–2682.
14 15	677	45. Nikolaidou ME, Marzilger SB, Mersmann F, Arampatzis A. Operating length and velocity
16 17 18	678	of human M. vastus lateralis fascicles during vertical jumping. R Soc Open Sci
19 20	679	2017;4:170185.
21 22 23 24	680	
25 26 27	681	
28 29 30	682	
31 32 33	683	
34 35 36	684	
37 38 39	685	
40 41 42	686	
43 44 45	687	
46 47 48	688	
49 50	689	
52 53	690	
55 56 57	691	
58 59 60	692	

TABLE AND FIGURE LEGENDS

Table 1. Study design timeline. Acronyms: CMJ = countermovement vertical jump; FWR = free-weight resistance; VR = variable resistance.

Table 2. Kinetic measures of vertical jump performance across all time points following the free-weight resistance and elastic band warm-up conditions (values are reported as mean \pm SE; *p < 0.05 compared to pre-intervention and FWR condition). Acronyms: Pre = pre-intervention; FWR = free-weight resistance; VR = variable resistance; K_{vert} = vertical stiffness; RFD = rate of force development.

Table 3. Kinematic measures of vertical jump performance across all time points following the free-weight resistance and elastic band warm-up conditions (values are reported as mean \pm SE; *p < 0.05 compared to pre-intervention). Acronyms: Pre = pre-intervention; FWR = free-weight resistance; VR = variable resistance; ECC = eccentric; CON = concentric.

Table 4. Normalized mean and peak VL, VM and Glut EMG amplitudes measured during vertical jumps across all time points following free-weight resistance and elastic band warm-up squat conditions (values are reported as mean \pm SE; *p < 0.05 compared to pre-intervention and FWR condition). Acronyms: VL = vastus lateralis; VM = vastus medialis; Glut = gluteus maximum; EMG = electromyogram; MVC = maximum voluntary contraction; FWR = free-weight resistance; VR = variable resistance; ECC = eccentric; CON = concentric.

Figure 1. Exemplar data from a subject depicting CMJ height, ground reaction force, knee angular velocity, knee flexion angle and VL EMG activity at 8 min following the free-weight resistance and elastic band warm-up squat conditions. VL = vastus lateralis; CMJ = countermovement vertical jump; FWR = free-weight resistance; VR = variable resistance.

Figure 2. Mean countermovement vertical jump height following free-weight resistance and elastic band warm-up squat conditions. *Significant increases (5.3-6.5%; p < 0.05) in vertical jump performance were achieved across all time points following the VR warm-up condition compared to pre-intervention and the FWR warm-up condition. Pre = pre-intervention; FWR = free-weight resistance; VR = variable resistance.

Table 1. Study design timeline.

Task	Time (min)
5-min cycle	0-5.0
5 unloaded squats (1 s/ 1 s)	5.0-6.0
5 unloaded squats (2 s/ 2 s)	6.0-7.0
5 CMJs (70%)	7.5-8.5
Single CMJs every 30 s (100%)	9.0-11.0
CMJ Test 1	13.0-13-5
FWR or VR squats	14.5-15.0
CMJ Tests (2-5)	15.5, 19.5, 23.5, 27.5
Jump; F W K = free-weight r variable resistance.	esistance; VR =

Table 2. Kinetic measures of vertical jump performance across all time points following the free-weight and variable resistance warm-up conditions (values are reported as mean \pm SE; *p < 0.05 compared to pre-intervention and FWR condition).

Measure	Condition	Pre	30 s	4 min	8 min	12 min
Peak Power (W)	FWR	49.3 ± 1.9	50.3 ± 1.5	50.8 ± 1.7	49.2 ± 2.2	50.3 ± 1.7
	VR	49.0 ± 1.7	51.3 ± 1.8*	51.8 ± 1.6*	51.0 ± 1.7*	51.2 ± 1.7*
Peak Eccentric Kinetic Energy (J)	FWR	87.4 ± 7.7	95.8 ± 9.0	90.8 ± 9.4	90.3 ± 7.7	88.0 ± 9.0
	VR	94.3 ± 8.0	93.6 ± 7.3	87.9 ± 7.2	88.8 ± 6.9	82.4 ± 5.9
Impulse asymmetry index (N·s)	FWR	2.9 ± 0.1	2.8 ± 0.1	2.9 ± 0.1	2.9 ± 0.1	2.9 ± 0.1
	VR	2.8 ± 0.1	2.9 ± 0.1	2.9 ± 0.1	2.9 ± 0.1	2.9 ± 0.1
Time asymmetry index (ms)	FWR	1.5 ± 0.1	1.4 ± 0.2	1.5 ± 0.2	1.3 ± 0.2	1.2 ± 0.2
	VR	1.4 ± 0.2	1.5 ± 0.2	1.1 ± 0.2	1.4 ± 0.2	1.2 ± 0.2
$K_{vert}(N^{-1}\cdot kg^{-1})$	FWR	70.8 ± 6.0	72.6 ± 3.8	70.5 ± 3.6	69.0 ± 3.7	72.0 ± 3.8
	VR	69.9 ± 5.0	73.3 ± 3.9	73.5 ± 4.5	74.6 ± 3.7	74.4 ± 4.5
Peak normalized RFD $(N \cdot s^{-1})$	FWR	134.2 ± 11.3	147.1 ± 12.2	132.5 ± 10.8	141.8 ± 13.5	118.2 ± 7.5
	VR	126.1 ± 6.7	$149.8 \pm 12.8*$	$143.2 \pm 11.7*$	$149.2 \pm 9.0*$	$147.7 \pm 13.7*$

Acronyms: Pre = pre-intervention; FWR = free-weight resistance; VR = variable resistance; K_{vert} = vertical stiffness; RFD = rate of force development.

Table 3. Kinematic measures of vertical jump performance across all time points following the free-weight
and variable resistance warm-up conditions (values are reported as mean \pm SE; * $p < 0.05$ compared to pre-
intervention).

Measure	Mode	Condition	Pre	30 s	4 min	8 min	12 min
Peak hip angular velocity $(^{\circ} \cdot s^{-1})$	ECC	FWR	301.1 ± 9.5	302.2 ± 10.0	294.4 ± 9.8	291.4 ± 6.9	292.8 ± 9.2
		VR	298.2 ± 7.1	305.2 ± 8.3	300.2 ± 8.3	302.6 ± 8.8	297.3 ± 8.7
	CON	FWR	584.6 ± 15.6	605.4 ± 18.8	591.9 ± 20.2	575.7 ± 20.7	576.2 ± 15.8
		VR	572.2 ± 17.1	591.9 ± 20.2	593.0 ± 22.1	588.2 ± 19.3	580.8 ± 21.7
Peak knee angular velocity ($^{\circ}\cdot s^{-1}$)	ECC	FWR	343.2 ± 13.6	341.0 ± 11.4	332.6 ± 11.9	343.8 ± 13.7	340.5 ± 14.3
• • •		VR	352.1 ± 14.3	364.9 ± 15.7	353.5 ± 13.9	363.0 ± 15.0	347.5 ± 16.6
	CON	FWR	956.4 ± 23.6	971.6 ± 24.6	969.3 ± 26.7	939.6 ± 27.9	959.6 ± 25.1
		VR	937.0 ± 23.8	966.0±28.8*	975.7 ± 29.7*	966.9 ± 26.2*	964.2 ± 24.5*
Peak ankle angular velocity ($^{\circ}\cdot s^{-1}$)	ECC	FWR	108.1 ± 10.0	117.6 ± 12.0	109.7 ± 12.2	112.0 ± 10.9	114.4 ± 10.8
		VR	121.1 ± 12.8	118.7 ± 9.8	120.3 ± 9.0	112.8 ± 7.1	104.5 ± 5.0
	CON	FWR	745.4 ± 23.4	733.7 ± 25.5	728.2 ± 18.9	707.9 ± 25.3	721.5 ± 27.0
		VR	717.9 ± 21.3	723.6 ± 22.2	731.7 ± 23.6	735.1 ± 28.1	739.4 ± 19.1
Peak hip flexion angle (°)		FWR	79.3 ± 2.0	82.7 ± 2.1	81.8 ± 1.7	82.1 ± 2.4	81.8 ± 2.2
		VR	81.5 ± 1.9	83.2 ± 1.4	83.3 ± 1.9	83.4 ± 1.5	82.8 ± 1.5
Peak knee flexion angle (°)		FWR	71.7 ± 2.9	73.3 ± 3.0	71.1 ± 2.8	72.0 ± 2.7	71.9 ± 3.3
		VR	71.8 ± 3.5	72.6 ± 3.4	74.2 ± 3.3	75.2 ± 2.8	75.4 ± 3.5
Peak ankle flexion angle (°)		FWR	32.7 ± 1.6	32.6 ± 1.4	32.7 ± 1.4	33.0 ± 1.4	33.2 ± 1.4
		VR	33.8 ± 1.6	34.7 ± 1.6	34.8 ± 1.6	35.2 ± 1.6	34.8 ± 1.8

Acronyms: Pre = pre-intervention; FWR = free-weight resistance; VR = variable resistance; ECC = eccentric; CON = concentric.

Table 4. Normalized mean and peak VL, VM and Glut EMG amplitudes measured during vertical jumps across all time points following free-weight and variable resistance warm-up squat conditions (values are reported as mean \pm SE; *p < 0.05 compared to pre-intervention and FWR condition).

Measure	Mode	Condition	Pre	30 s	4 min	8 min	12 min
Mean VL EMG (%MVC)	ECC	FWR	31.9 ± 2.1	31.5 ± 3.4	30.5 ± 3.0	30.0 ± 2.0	29.5 ±2.9
		VR	28.5 ± 2.1	30.2 ± 2.2	31.9 ± 2.3	31.5 ± 1.8	32.0 ± 2.2
	CON	FWR	82.4 ± 6.1	83.9 ± 6.4	78.4 ± 8.5	72.5 ± 4.6	73.4 ± 6.0
		VR	85.4 ± 7.8	$108.2 \pm 13.3^*$	$110.2 \pm 12.2*$	$107.5 \pm 10.5*$	102.6 ± 9.4*
Peak VL EMG (%MVC)	ECC	FWR	89.2 ± 6.7	94.7 ± 8.0	90.4 ± 6.9	85.0 ± 6.7	84.0 ± 6.0
		VR	91.6 ± 4.9	95.0 ± 8.0	94.2 ± 6.1	90.7 ± 4.7	90.6 ± 5.9
	CON	FWR	112.4 ± 8.0	123.7 ± 9.6	116.9 ± 7.1	112.8 ± 6.7	112.2 ± 6.3
		VR	114.3 ± 5.8	115.8 ± 10.2	117.4 ± 7.6	111.2 ± 6.6	108.1 ± 6.6
Mean VM EMG (%MVC)	ECC	FWR	36.3 ± 2.7	33.9 ± 4.1	32.6 ± 3.3	32.2 ± 2.6	33.0 ± 3.4
. ,		VR	37.9 ± 3.8	40.2 ± 3.8	38.1 ± 3.3	39.9 ± 2.9	38.1 ± 3.3
	CON	FWR	94.9 ± 5.0	95.3 ± 10.9	85.1 ± 8.4	85.1 ± 6.4	87.7 ± 7.3
		VR	90.2 ± 9.2	96.0 ± 7.9	94.7 ± 7.5	88.8 ± 5.9	87.2 ± 6.2
Peak VM EMG (%MVC)	ECC	FWR	98.6 ± 7.9	96.9 ± 7.4	92.9 ± 6.7	88.0 ± 6.1	89.0 ± 6.1
		VR	111.3 ± 9.5	114.4 ± 11.2	108.1 ± 10.2	104.7 ± 8.1	97.3 ± 6.7
	CON	FWR	132.0 ± 12.2	128.7 ± 11.5	120.5 ± 9.5	118.2 ± 8.6	118.8 ± 9.1
		VR	150.6 ± 12.8	149.0 ± 14.4	143.4 ± 12.5	140.8 ± 11.8	127.7 ± 8.9
Mean Glut EMG (%MVC)	ECC	FWR	20.0 ± 1.9	22.4 ± 2.8	21.3 ± 2.3	19.7 ± 1.7	21.2 ± 1.9
		VR	21.7 ± 2.5	23.2 ± 2.6	22.5 ± 2.4	22.5 ± 2.2	21.7 ± 2.0
	CON	FWR	81.1 ± 10.2	84.4 ± 14.8	83.9 ± 12.1	78.6 ± 8.7	75.1 ± 9.2
		VR	84.4 ± 14.0	87.1 ± 14.2	78.3 ± 10.1	77.4 ± 8.5	76.8 ± 9.8
Peak Glut EMG (%MVC)	ECC	FWR	72.6 ± 7.0	78.3 ± 7.5	76.4 ± 7.2	74.6 ± 7.1	71.7 ± 6.9
		VR	79.1 ± 9.4	82.0 ± 9.5	76.7 ± 8.9	74.7 ± 5.5	71.7 ± 7.3
	CON	FWR	103.4 ± 10.8	111.3 ± 14.0	112.7 ± 12.5	103.3 ± 12.1	103.1 ± 9.6
		VR	115.9 ± 9.9	118.5 ± 10.7	110.8 ± 11.7	100.3 ± 7.9	99.7 ± 7.6

Acronyms: VL = vastus lateralis; VM = vastus medialis; Glut = gluteus maximum; EMG = electromyogram; MVC = maximum voluntary contraction; FWR = free-weight resistance; VR = variable resistance; ECC = eccentric; CON = concentric.





300x500mm (96 x 96 DPI)



60





250x200mm (96 x 96 DPI)